

**Atomic structure of a toxic, oligomeric segment of SOD1 linked to amyotrophic lateral sclerosis (ALS).**

**Journal:** Proc Natl Acad Sci U S A

**Publication Year:** 2017

**Authors:** Smriti Sangwan, Anni Zhao, Katrina L Adams, Christina K Jayson, Michael R Sawaya, Elizabeth L Guenther, Albert C Pan, Jennifer Ngo, Destaye M Moore, Angela B Soriaga, Thanh D Do, Lukasz Goldschmidt, Rebecca Nelson, Michael T Bowers, Carla M Koehler, David E Shaw, Bennett G Novitch, David S Eisenberg

**PubMed link:** 28760994

**Funding Grants:** Molecular Characterization of hESC and iPSC-Derived Spinal Motor Neurons , CSUN-UCLA Bridges to Stem Cell Research, In vitro modeling of human motor neuron disease , A small molecule tool for reducing the malignant potential in reprogramming human iPSCs and ESCs

**Public Summary:**

More than 170 mutations in superoxide dismutase 1 (SOD1) are linked to inherited forms of ALS, and aggregates of this protein are a pathological feature associated with this disease. Although it is accepted that SOD1 gains a toxic function in the disease state, a molecular understanding of the toxic species is lacking. Here, we identify a short segment of SOD1 that is both necessary and sufficient for toxicity to motor neurons. The crystal structure of the segment reveals an out-of-register  $\beta$ -sheet oligomer, providing a structural rationale for the toxic effects of mutant SOD1 in ALS.

**Scientific Abstract:**

Fibrils and oligomers are the aggregated protein agents of neuronal dysfunction in ALS diseases. Whereas we now know much about fibril architecture, atomic structures of disease-related oligomers have eluded determination. Here, we determine the corkscrew-like structure of a cytotoxic segment of superoxide dismutase 1 (SOD1) in its oligomeric state. Mutations that prevent formation of this structure eliminate cytotoxicity of the segment in isolation as well as cytotoxicity of the ALS-linked mutants of SOD1 in primary motor neurons and in a *Danio rerio* (zebrafish) model of ALS. Cytotoxicity assays suggest that toxicity is a property of soluble oligomers, and not large insoluble aggregates. Our work adds to evidence that the toxic oligomeric entities in protein aggregation diseases contain antiparallel, out-of-register beta-sheet structures and identifies a target for structure-based therapeutics in ALS.

---

**Source URL:** <https://www.cirm.ca.gov/about-cirm/publications/atomic-structure-toxic-oligomeric-segment-sod1-linked-amyotrophic-lateral>