
Comparison of Magnetic Resonance Imaging and Serum Biomarkers for Detection of Human Pluripotent Stem Cell-Derived Teratomas.

Journal: Stem Cell Reports

Publication Year: 2016

Authors: Johannes Riegler, Antje Ebert, Xulei Qin, Qi Shen, Mouer Wang, Mohamed Ameen, Kazuki Kodo, Sang-Ging Ong, Won Hee Lee, Grace Lee, Evgenios Neofytou, Joseph D Gold, Andrew J Connolly, Joseph C Wu

PubMed link: 26777057

Funding Grants: Human Embryonic Stem Cell-Derived Cardiomyocytes for Patients with End Stage Heart Failure, Macaca mulatta as advanced model for predictive preclinical testing of engineered cardiac autografts and allografts

Public Summary:

The use of cells derived from pluripotent stem cells (PSCs) for regenerative therapies confers a considerable risk for neoplastic growth and teratoma formation. Preclinical and clinical assessment of such therapies will require suitable monitoring strategies to understand and mitigate these risks. Here we generated human-induced pluripotent stem cells (iPSCs), selected clones that continued to express reprogramming factors after differentiation into cardiomyocytes, and transplanted these cardiomyocytes into immunocompromised rat hearts post-myocardial infarction. We compared magnetic resonance imaging (MRI), cardiac ultrasound, and serum biomarkers for their ability to delineate teratoma formation and growth. MRI enabled the detection of teratomas with a volume >8 mm³. A combination of three plasma biomarkers (CEA, AFP, and HCG) was able to detect teratomas with a volume >17 mm³ and with a sensitivity of more than 87%. Based on our findings, a combination of serum biomarkers with MRI screening may offer the highest sensitivity for teratoma detection and tracking.

Scientific Abstract:

The use of cells derived from pluripotent stem cells (PSCs) for regenerative therapies confers a considerable risk for neoplastic growth and teratoma formation. Preclinical and clinical assessment of such therapies will require suitable monitoring strategies to understand and mitigate these risks. Here we generated human-induced pluripotent stem cells (iPSCs), selected clones that continued to express reprogramming factors after differentiation into cardiomyocytes, and transplanted these cardiomyocytes into immunocompromised rat hearts post-myocardial infarction. We compared magnetic resonance imaging (MRI), cardiac ultrasound, and serum biomarkers for their ability to delineate teratoma formation and growth. MRI enabled the detection of teratomas with a volume >8 mm³. A combination of three plasma biomarkers (CEA, AFP, and HCG) was able to detect teratomas with a volume >17 mm³ and with a sensitivity of more than 87%. Based on our findings, a combination of serum biomarkers with MRI screening may offer the highest sensitivity for teratoma detection and tracking.

Source URL: <https://www.cirm.ca.gov/about-cirm/publications/comparison-magnetic-resonance-imaging-and-serum-biomarkers-detection-human>