

Distinct and separable roles for EZH2 in neurogenic astroglia.

Journal: Elife

Publication Year: 2014

Authors: William W Hwang, Ryan D Salinas, Jason J Siu, Kevin W Kelley, Ryan N Delgado, Mercedes F Paredes, Arturo Alvarez-Buylla, Michael C Oldham, Daniel A Lim

PubMed link: 24867641

Funding Grants: Training Program in Stem Cell Research at UCSF

Public Summary:

The epigenetic mechanisms that enable specialized astrocytes to retain neurogenic competence throughout adult life are still poorly understood. Here we show that astrocytes that serve as neural stem cells (NSCs) in the adult mouse subventricular zone (SVZ) express the histone methyltransferase EZH2. This Polycomb repressive factor is required for neurogenesis independent of its role in SVZ NSC proliferation, as *Ink4a/Arf*-deficiency in *Ezh2*-deleted SVZ NSCs rescues cell proliferation, but neurogenesis remains defective. *Olig2* is a direct target of EZH2, and repression of this bHLH transcription factor is critical for neuronal differentiation. Furthermore, *Ezh2* prevents the inappropriate activation of genes associated with non-SVZ neuronal subtypes. In the human brain, SVZ cells including local astroglia also express EZH2, correlating with postnatal neurogenesis. Thus, EZH2 is an epigenetic regulator that distinguishes neurogenic SVZ astrocytes, orchestrating distinct and separable aspects of adult stem cell biology, which has important implications for regenerative medicine and oncogenesis. DOI: <http://dx.doi.org/10.7554/eLife.02439.001>.

Scientific Abstract:

The epigenetic mechanisms that enable specialized astrocytes to retain neurogenic competence throughout adult life are still poorly understood. Here we show that astrocytes that serve as neural stem cells (NSCs) in the adult mouse subventricular zone (SVZ) express the histone methyltransferase EZH2. This Polycomb repressive factor is required for neurogenesis independent of its role in SVZ NSC proliferation, as *Ink4a/Arf*-deficiency in *Ezh2*-deleted SVZ NSCs rescues cell proliferation, but neurogenesis remains defective. *Olig2* is a direct target of EZH2, and repression of this bHLH transcription factor is critical for neuronal differentiation. Furthermore, *Ezh2* prevents the inappropriate activation of genes associated with non-SVZ neuronal subtypes. In the human brain, SVZ cells including local astroglia also express EZH2, correlating with postnatal neurogenesis. Thus, EZH2 is an epigenetic regulator that distinguishes neurogenic SVZ astrocytes, orchestrating distinct and separable aspects of adult stem cell biology, which has important implications for regenerative medicine and oncogenesis. DOI: <http://dx.doi.org/10.7554/eLife.02439.001>.

Source URL: <https://www.cirm.ca.gov/about-cirm/publications/distinct-and-separable-roles-ezh2-neurogenic-astroglia>