

---

**Imaging dynamic and selective low-complexity domain interactions that control gene transcription.**

<b>Journal:</b>	Science
<b>Publication Year:</b>	2018
<b>Authors:</b>	Shasha Chong, Claire Dugast-Darzacq, Zhe Liu, Peng Dong, Gina M Dailey, Claudia Cattoglio, Alec Heckert, Sambashiva Banala, Luke Lavis, Xavier Darzacq, Robert Tjian
<b>PubMed link:</b>	29930090
<b>Funding Grants:</b>	Single Molecule Biophysics and Biology of Cellular Identity

**Public Summary:**

Many eukaryotic transcription factors (TFs) contain intrinsically disordered low-complexity sequence domains (LCDs), but how these LCDs drive transactivation remains unclear. We used live-cell single-molecule imaging to reveal that TF LCDs form local high-concentration interaction hubs at synthetic and endogenous genomic loci. TF LCD hubs stabilize DNA binding, recruit RNA polymerase II (RNA Pol II), and activate transcription. LCD-LCD interactions within hubs are highly dynamic, display selectivity with binding partners, and are differentially sensitive to disruption by hexanediols. Under physiological conditions, rapid and reversible LCD-LCD interactions occur between TFs and the RNA Pol II machinery without detectable phase separation. Our findings reveal fundamental mechanisms underpinning transcriptional control and suggest a framework for developing single-molecule imaging screens for drugs targeting gene regulatory interactions implicated in disease.

**Scientific Abstract:**

Many eukaryotic transcription factors (TFs) contain intrinsically disordered low-complexity sequence domains (LCDs), but how these LCDs drive transactivation remains unclear. We used live-cell single-molecule imaging to reveal that TF LCDs form local high-concentration interaction hubs at synthetic and endogenous genomic loci. TF LCD hubs stabilize DNA binding, recruit RNA polymerase II (RNA Pol II), and activate transcription. LCD-LCD interactions within hubs are highly dynamic, display selectivity with binding partners, and are differentially sensitive to disruption by hexanediols. Under physiological conditions, rapid and reversible LCD-LCD interactions occur between TFs and the RNA Pol II machinery without detectable phase separation. Our findings reveal fundamental mechanisms underpinning transcriptional control and suggest a framework for developing single-molecule imaging screens for drugs targeting gene regulatory interactions implicated in disease.

---

**Source URL:** <https://www.cirm.ca.gov/about-cirm/publications/imaging-dynamic-and-selective-low-complexity-domain-interactions-control>