

Recurrent variations in DNA methylation in human pluripotent stem cells and their differentiated derivatives.

Journal: Cell Stem Cell

Publication Year: 2012

Authors: Kristopher L Nazor, Gulsah Altun, Candace Lynch, Ha Tran, Julie V Harness, Ileana Slavin, Ibon Garitaonandia, Franz-Josef Muller, Yu-Chieh Wang, Francesca S Boscolo, Eyitayo Fakunle, Biljana Dumevska, Sunray Lee, Hyun Sook Park, Tsaiwei Olee, Darryl D D'Lima, Ruslan Semechkin, Mana M Parast, Vasily Galat, Andrew L Laslett, Uli Schmidt, Hans S Keirstead, Jeanne F Loring, Louise C Laurent

PubMed link: 22560082

Funding Grants: Molecular Mechanisms of Trophoblast Stem Cell Specification and Self-Renewal , TSRI Center for hESC Research, The Stem Cell Matrix: a map of the molecular pathways that define pluripotent cells, Ensuring the safety of cell therapy: a quality control pipeline for cell purification and validation, Thymus based tolerance to stem cell therapies , Collaborative Laboratory for Human Embryonic Stem Cell Research at Sanford-Burnham Medical Research Institute

Public Summary:

We report the most comprehensive study of human pluripotent stem cell variation to date. We identified recurrent aberrations in X chromosome inactivation and imprinting that were maintained during differentiation. DNA hypomethylation was the most discriminate epigenetic feature of any tissue and was recapitulated by DNA demethylation during in vitro directed differentiation.

Scientific Abstract:

Human pluripotent stem cells (hPSCs) are potential sources of cells for modeling disease and development, drug discovery, and regenerative medicine. However, it is important to identify factors that may impact the utility of hPSCs for these applications. In an unbiased analysis of 205 hPSC and 130 somatic samples, we identified hPSC-specific epigenetic and transcriptional aberrations in genes subject to X chromosome inactivation (XCI) and genomic imprinting, which were not corrected during directed differentiation. We also found that specific tissue types were distinguished by unique patterns of DNA hypomethylation, which were recapitulated by DNA demethylation during in vitro directed differentiation. Our results suggest that verification of baseline epigenetic status is critical for hPSC-based disease models in which the observed phenotype depends on proper XCI or imprinting and that tissue-specific DNA methylation patterns can be accurately modeled during directed differentiation of hPSCs, even in the presence of variations in XCI or imprinting.

Source URL: <https://www.cirm.ca.gov/about-cirm/publications/recurrent-variations-dna-methylation-human-pluripotent-stem-cells-and-their>